

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 921 192 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
09.06.1999 Bulletin 1999/23

(21) Application number: 97203781.6

(22) Date of filing: 03.12.1997

(51) Int. Cl.⁶: **C12N 15/12, C07K 14/47,**
A61K 38/17, C12Q 1/68,
C12N 1/19, C12N 1/21

(84) Designated Contracting States:
AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(71) Applicant: Leadd B.V.
2333 AL Leiden (NL)

(72) Inventors:
• Noteborn, Mathieu Hubertus Maria
2352 EH Lelidorp (NL)

• Danen - van Oorschot, Astrid A. A. M.
2651 VG Berkel en Rodenrijs (NL)

(74) Representative:
Smulders, Theodorus A.H.J., Ir.
Vereenigde Octroolbureaux
Nieuwe Parklaan 97
2587 BN 's-Gravenhage (NL)

Remarks:

The applicant has subsequently filed a sequence
listing and declared, that it includes no new matter.

(54) Molecules interacting with apoptin

(57) The invention relates to activation of apoptosis
by means of interference of Hou-like and/or IFP35-like
compounds.

Also the invention relates to anti-tumor therapies
with compounds, which negatively interfere with Hou-
like and/or IFP35-like compounds leading to induction
of apoptosis, resulting in the elimination of tumor cells.

Also the invention relates to therapies for diseases
related to aberrant apoptosis induction, such as auto-
immune disease.

Also the invention describes the diagnosis of cells,
which are susceptible to apoptin- or apoptin-like
induced apoptosis.

EP 0 921 192 A1

new treatments and diagnosis for diseases related with aberrancies in the apoptotic process, such as cancer and autoimmune diseases.

[0013] Proteins found associating with apoptin include members of the family of Nmi/Hou-like and IFP-like proteins.

[0014] Thus the invention provides a recombinant and/or isolated nucleic acid molecule encoding at least a functional part of a member of the family of Nmi-like proteins or at least a functional part of a member of the family of Hou-like proteins or at least a functional part of a member of the family of IFP35-like proteins for use in the induction of apoptosis in a population of cells related to a pathological condition.

[0015] As explained herein the expression of Hou is connected to oncogenes and has been found to be high in certain transformed cells. These are typically the cells that can be induced to go into apoptosis by apoptotic agents such as apoptin. Typically providing a cell with Hou-like activity will therefor increase the chance of inducing apoptosis in such a cell. IFP35-like proteins are involved in transporting apoptotic substances to the nucleus of cells. Under influence of for instance interferons these proteins localize in the nucleus. Therefor IFP-like activity is used to get apoptin-like activity into the nucleus, which is important for the induction of apoptosis, for instance through Hou-like proteins. The Hou-like activity or Nmi-like activity is defined herein as any molecule capable of exerting the same or a similar function as the original Hou-like (Nmi-like) protein. The same definition goes for IFP-activity. Typically such a molecule can be encoded by a nucleic acid molecule which comprises at least a functional and specific part of the sequence of figure 1, 2, 4 or 5 or encoding an amino sequence of figure 6 or a sequence at least 60, preferably 70, preferably 90 % homologous with said functional and specific sequence or comprising a sequence hybridizing to any of the foregoing sequences under stringent conditions. In order to be able to express the Hou-like activity and/or the IFP-like activity it is preferred to have an expression vector encoding said activity. Expression vectors are nucleic acid molecules which can be brought into cells, or transfect cells themselves and which have the machinery (together with the machinery of the host cell) to express proteins encoded on the expression vector when present in a cell.

[0016] It is preferred that cells which are provided, according to the invention, with Hou-like activity and/or IFP-like activity, are also provided with apoptosis inducing activity, preferably apoptin-like activity, which is defined along the same lines as Hou-like activity. In order to get the activity into the cells in which apoptosis has to be induced it is possible and preferred to use a gene delivery vehicle. A gene delivery vehicle is a means to transport a nucleic acid molecule capable of expressing the wanted activity in a host cell into said host cell. Gene delivery vehicles are known in the art. They include for instance recombinant viruses such as adenoviruses and retroviruses, but also non-viral vehicles such as polymers and liposomes have been suggested. Methods of targeting gene delivery vehicles to target cells are also known in the art and need not be elaborated herein. The invention also provides the newly identified molecules themselves, both the nucleic acid molecules (meaning DNA coding and/or non coding strands as well as RNA) and the proteinaceous molecules (peptides, polypeptides, glycoproteins and associations between proteins and RNA's and the like). Based on the given sequences other family members of the Hou/Nmi and IFP families will be identified having the same or similar function. Typically such molecules will have high homology to the sequences given herein.

[0017] For nucleic acid molecules the homology is expected to be at least 60, preferably 70, more preferably 80% therewith.

[0018] These nucleic acid molecules can of course again be incorporated into expression vectors as mentioned hereinbefore. Preferably these expression vectors also encode apoptotic activity, preferably apoptin or a functional fragment and/or equivalent thereof.

[0019] These expression vectors can again be made into gene delivery vehicles.

[0020] The invention also provides the recombinant or isolated proteinaceous substance comprising at least a functional part of a member of the family of Nmi/Hou-like proteins or at least a functional part of a member of the family of Hou-like proteins for use in the induction of apoptosis in a population of cells related to a pathological condition and an Nmi/Hou-like proteinaceous substance having at least a functional and/or specific part of the sequence of figure 3 or being encoded by a functional and/or specific part of the sequence of figure 1 or figure 2 or being at least 60, preferably 70, preferably 80% homologous to at least a functional and/or specific part of the sequence of figure 3 or being at least 60, preferably 70, preferably 80% homologous to a protein encoded by at least a functional and/or specific part of the sequence of figure 1 or figure 2 and an IFP35-like proteinaceous substance having at least a functional and/or specific part of the sequence of figure 6 or 7 or being encoded by a functional and/or specific part of the sequence of figure 4 or figure 5 or being at least 60, preferably 70, preferably 80% homologous to at least a functional and/or specific part of the sequence of figure 6 or 7 or being at least 60, preferably 70, preferably 80% homologous to a protein encoded by at least a functional and/or specific part of the sequence of figure 4 or figure 5.

[0021] A functional part in this document means having the same or similar activity (although the amount of activity may differ) A specific part herein means a part of sufficient size to be specific for the protein or nucleic acid or to be of sufficient size to distinguish the protein from another protein immunologically. The proteins disclosed herein can for instance also be used to identify further components of the apoptotic pathway.

[0022] The reason for bringing IFP-like activity and/or Hou-like activity together with apoptotic activity is of course to induce aberrant cells to go into apoptosis. Thus the invention also provides a method for inducing apoptosis in cells

GAL4-activation domain-tagged cDNA library

[0043] The expression vector pACT, containing the cDNAs from Epstein-Barr-virus-transformed human B cells fused to the GAL4 transcriptional activation domain, was used for detecting apoptin-associating proteins. The pACT c-DNA library is derived from the lambda-ACT cDNA library, as described by Durfee et al. 1993.

Bacterial and Yeast strains

[0044] The E.coli strain JM109 was the transformation recipient for the plasmid pGBT9 and pGBT-VP3. The bacterial strain electromax/DH10B was used for the transformation needed for the recovery the apoptin-associating pACT-cDNAs, and was obtained from GIBCO-BRL, USA.

[0045] The yeast strain Y190 was used for screening the cDNA library, and all other transformations which are part of the used yeast-two-hybrid system.

Media

[0046] For drug selections Luria Broth (LB) plates for E.coli were supplemented with ampicillin (50 microgram per ml). Yeast YPD and SC media were prepared as described by Rose et al. (1990).

Transformation of competent yeast strain Y190 with plasmids pGBT-VP3 and pACT-cDNA and screening for beta-galactosidase activity.

[0047] The yeast strain Y190 was made competent and transformed according to the methods described by Klebe et al. (Klebe et al., 1983). The yeast cells were first transformed with pGBT-VP3 and subsequently transformed with pACT-cDNA, and these transformed yeast cells were grown on histidine-minus plates, also lacking leucine and tryptophan.

[0048] Hybond-N filters were layed on yeast colonies, which were histidine-positive and allowed to wet completely. The filters were lifted and submerged in liquid nitrogen to permeabilize the yeast cells. The filters were thawed and layed with the colony side up on Whatman 3MM paper in a petridish with Z-buffer (Per liter: 16.1 gr $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 5.5 gr $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$, 0.75 gr KCl and 0.246 gr $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, pH 7.0) containing 0.27% beta-mercapto-ethanol and 1 mg/ml X-gal. The filters were incubated for at least 15 minutes or during night.

Recovery of plasmids from yeast

[0049] Total DNA from yeast cells, which were histidine- and beta-galactosidase-positive, was prepared by using the glucanase-alkaline lysis method as described by Hoffman and Winston (1987) and used to transform Electromax/DH10B bacteria via electroporation using a Bio-Rad GenePulser according to the manufacturer's specifications.

[0050] Transformants were plated on LB media containing ampicillin.

Isolation of apoptin-associating pACT clones

[0051] By means of colony-filter assay the colonies were lysed and hybridized to a radioactive-labeled 17-mer oligomer, which is specific for pACT (see also section Sequence analysis).

[0052] Plasmid DNA was isolated from the pACT-clones, and by means of *Xho*I digestion analysed for the presence of a cDNA insert.

Sequence analysis

[0053] The subclones containing the sequences encoding apoptin-associating proteins were sequenced using dideoxy NTPs according to the Sanger method which was performed by Eurogentec, Nederland BV (Maastricht, The Netherlands). The used sequencing primer was a pACT-specific 17-mer comprising of the DNA-sequence 5'-TACCACTACAATGGATG-3'.

[0054] The sequences of the apoptin-associating proteins were compared with known gene sequences from the EMBL/Genbank.

Results and discussion

[0055] Apoptin induces specifically apoptosis in transformed cells, such as cell lines derived from human tumors. To identify the essential compounds in this cell-transformation-specific and/or tumor-specific apoptosis pathway, a yeast

Nmi, or Hou will be interchangeably used.

[0070] In this respect, the pattern of Nmi expression is interesting, since it is expressed at low levels in normal tissues, in contrast to its high levels of expression in transformed cell lines. Among eight cancer lines tested, highest levels were observed in four leukemia cell lines (Ba and Zervos, 1996).

[0071] In leukemias, a high expression of C-myc correlates with a high level of Nmi (HL-60, K562 and MOLT-4). The Nmi gene is located on chromosome 22, which is also involved in the t (9;22) translocation leading to the Bcr-Abl fusion protein, as seen in some leukemias (Rabbits, 1991, Sawyers and Deny, 1994).

[0072] Using a yeast genetic screen, Nmi was identified as a protein that binds to N-myc and C-myc. Myc proteins are important in the regulation of cell proliferation and differentiation. Together with ras or raf, myc can transform primary cells in culture. Nmi/Hou-like proteins will up-regulate the activity of Myc proteins via binding to them.

[0073] Up-regulation of Myc proteins has been described for Burkitt lymphomas, neuroblastomas and small cell lung carcinomas. Myc proteins contain a basic region, a helix-loop-helix (HLH) and a leucine zipper (Zip), and form homo- or heterodimers that can bind to specific DNA sequences and regulate transcription. Myc also forms heterodimers with Max. Myc/Max heterodimers activate transcription, whereas Max homodimers repress transcription, thus antagonizing Myc's function (Evan and Littlewood, 1993).

[0074] Nmi was found to interact with N-myc, c-myc, Max, Mxi1 and other transcription factors that have HLH and/or Zip motifs. Interaction with N-myc and C-myc was confirmed by co-precipitation experiments (Bao and Zervos, 1996).

Induction of apoptosis through interference with the function of Nmi/Hou-like proteins.

[0075] Our results indicate that apoptin can change the Nmi/Hou-like-mediated proliferation (transformation/tumor-formation) activity into a Nmi/Hou-like-mediated apoptotic activity. Remarkably, this Nmi/Hou-like-mediated apoptotic activity will be specific for transformed/tumor cells, due to the very high level of Nmi/Hou in transformed cells in combination with over-expression of (proto-)oncogenes, such as Myc.

[0076] By means of transient transfection assays, it was shown that over-expression of the determined Hou-like protein (see Fig. 3) and apoptin did result in induction of apoptosis in normal VH10-, VH25-fibroblasts. In contrast to normal fibroblasts which over-expressed only apoptin. This result indicates that Hou-like proteins are an important factor in (apoptin-induced) apoptosis.

[0077] The presented data imply that interference with the function of Nmi/Hou-like proteins resulting in apoptosis can be used as a specific anti-tumor therapy, or therapies of related diseases, such as auto-immune diseases.

Characteristics of the apoptin-associating protein IFP35

[0078] The other apoptin-associating protein is IFP35, which is an interferon(IFN)-induced leucine zipper protein of 282 a.a., and has an apparent molecular mass of 35 kD. It was isolated by differential screening from HeLa cells that had been treated with IFN- γ (Bange et al., 1994).

[0079] IFP35 mRNA could be induced by IFN- γ in different human cell types, including fibroblasts, macrophages, and epithelial cells. It has a leucine zipper motif at the N-terminus, but it lacks an adjacent basic domain required for DNA binding. It has been suggested that these types of proteins negatively regulate bZIP transcription factors by forming non-functional heterodimers. IFP35 was shown to form homodimers (Bange et al., 1994).

Induction of apoptosis by interference of IFP35 in combination with Hou/Nmi-like proteins.

[0080] IFP35 is found in the cell nucleus, after interferon treatment and is expressed in a wide variety of cell types including fibroblasts, macrophages and epithelial cells (Bange et al., 1994).

[0081] In general, virus infections trigger interferon production. It is likely that a CAV infection and/or expression of apoptin will result in interferon up-regulation, which might result in the translocation of IFP35 or IFP35-like proteins into the nucleus. IFP35 will transport apoptin also to the nucleus, due to its association.

[0082] It seems likely that if apoptin is transported into the nucleus by IFP35 it will be able to associate with the IFP35-homologous region within Hou/Nmi-like proteins. This association will cause an aberrant regulation of Hou/Nmi-regulated genes, such as the oncogene Myc. Subsequently, the cells over-expressing Nmi/Hou-like proteins and oncogenes, such as Myc will undergo apoptosis.

[0083] Experimental evidence for IFP35 as an essential factor in (apoptin) apoptosis induction was derived from the following experiments. Normal VH10 cells over-expressing Hou/Nmi, IFP35 and apoptin underwent faster apoptosis than normal VH10 cells expressing Hou/Nmi and apoptin.

retinoblastoma protein associates with the protein phosphate type I catalytic subunit. *Genes and Development* 7, 555-569.

8. Earnshaw, W.C., 1995. Nuclear changes in apoptosis. *Current Opinion in Cell Biology* 7, 337-343.

9. Evan, G.I. and Littlewood, T.D. (1993). *Current Opinions in Genetics Development* 3, 44-49.

10. Fields, S. and Song, O.K. (1989). A novel genetic system to detect protein-protein interactions. *Nature* 340, 245-246.

11. Hockenberry, D.M. (1994). Bcl-2 in cancer, development and apoptosis. *Journal of Cell Science, Supplement* 18, 51-55.

12. Hoffman, C.S. and Winston, F. (1987). A ten-minute DNA preparation from yeast efficiently releases autonomous plasmids for transformation of *Escherichia coli*. *Gene* 57, 267-272.

13. Kerr, J.F.R., Winterford, C.M., and Harmon, B.V. (1994). Apoptosis: Its significance in cancer and cancer therapy. *Cancer* 73, 2013-2026.

14. Klebe, R.J., Harriss, J.V., Sharp, Z.D., and Douglas, M.G. (1983). A general method for polyethylene-glycol-induced genetic transformation of bacteria and yeast. *Gene* 25, 333-341.

15. Levine, A.J. (1997). p53, the cellular gatekeeper for growth and division. *Cell* 88, 323-331.

16. Maniatis, T., Fritsch, E.F., and Sambrook, J. (1982). *Molecular Cloning: A Laboratory Manual*. CSHL Press, New York, USA.

17. Matsudaira, P. (1994). Actin crosslinking proteins at the leading edge. *Seminars in Cell biology* 5, 165-174.

18. McDonnell T.J., Meyn, R.E., Robertson, L.E. (1995). Implications of apoptotic cell death regulation in cancer therapy. *Seminars in Cancer Biology* 6, 53-60.

19. Noteborn, M.H.M. (1996). PCT application WO 96/41191. Apoptin induces apoptosis in human transformed and malignant cells but not in normal cells as essential characteristic for the development of an anti-tumor therapy.

20. Noteborn, M.H.M., and De Boer, G.F. (1996). Patent USA/no. 030, 335.

21. Noteborn, M.H.M., De Boer, G.F., Van Roozelaar, D., Karreman, C., Kranenburg, O., Vos, J., Jeurissen, S., Zanterra, A., Hoeber, R., Koch, G., Van Ormondt, H., and Van der Eb, A.J. (1991). Characterization of cloned chicken anemia virus DNA that contains all elements for the infectious replication cycle. *Journal of Virology* 65, 3131-3139.

22. Noteborn, M.H.M., Hoeber, R.C., and Pietersen, A. (1997). A gene delivery vehicle expressing the apoptosis-inducing proteins VP2 and/or apoptin. European Patent Application no. 97201121.7

23. Noteborn, M.H.M., Todd, D., Verschueren, C.A.J., De Gauw, H.W.F.M., Curran, W.L., Veldkamp, S., Douglas, A.J., McNulty, M.S., Van der Eb, A.J., and Koch, G. (1994). A single chicken anemia virus protein induces apoptosis. *Journal of Virology* 68, 346-351.

24. Noteborn, M.H.M., and Zhang, Y. (1997). Methods and means for determining the transforming capability of agents, for determining the predisposition of cells to become transformed and prophylactic treatment of cancer using apoptin-like activity. European Patent Application no. 97439

25. Paulovich, A.G., Toczyski, D., Hartwell, H. (1997). When checkpoints fail. *Cell* 88, 315-321.

26. Rabbitts, T.H. (1991). *Cell* 67, 641-644.

27. Rose, M.D., Winston, F., and Hieter, P. (1990). *Methods in yeast genetics. A laboratory course manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, USA.

28. Sachs, L. and Lotem, J. (1993). Control of programmed cell death in normal and leukemia cells: New implications for therapy. *Blood* 82, 15-21.

29. Sanger, F., Nicklen, S., and Coulson, A.R. (1977). DNA sequencing with chain-terminating inhibitors. *Proceedings National Academic Sciences USA* 74, 5463-5467.

30. Steller, H. (1995). Mechanisms and genes of cellular suicide. *Science* 267, 1445-1449.

31. Sawyers, C.L. and Denny, C.T. (1994) *Cell* 77, 171-173.

32. Telford, W.G., King, L.E., Fraker, P.J. (1992). Comparative evaluation of several DNA binding dyes in the detection of apoptosis-associated chromatin degradation by flow cytometry. *Cytometry* 13, 137-143.

33. Teodoro, J.G. and Branton, P.E. (1997). Regulation of apoptosis by viral gene products. *Journal of Virology* 71, 1739-1746.

34. Thompson, C.B. (1995). Apoptosis in the pathogenesis and treatment of disease. *Science* 267, 1456-1462.

35. White, E. (1996). Life, death, and the pursuit of apoptosis. *Genes and development* 10, 1-15.

36. Wyllie, A.H. (1995). The genetic regulation of apoptosis. *Current Opinion in Genetics and Development* 5, 97-104.

37. Wyllie, A.H., Kerr, J.F.R., Currie, A.R. (1980). Cell death: The significance of apoptosis. *International Review of Cytology* 68, 251-306.

38. Yang, X., Hubbard, E.J.A., and Carlson, M. (1992). A protein kinase substrate identified by the two-hybrid system. *Science* 257, 680-682.

39. Zhuang, S.-M., Landegent, J.E., Verschueren, C.A.J., Falkenburg, J.H.F., Van Ormondt, H., Van der Eb, A.J.,

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: Leadd B.V.
- (B) STREET: Wassenaarseweg 72
- (C) CITY: Leiden
- (D) STATE: Zuid-Holland
- (E) COUNTRY: the Netherlands
- (F) POSTAL CODE (ZIP): 2333 AL

(ii) TITLE OF INVENTION: Novel molecules involved in
apoptotic pathways.

(iii) NUMBER OF SEQUENCES: 14

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version

#1.30 (EPO)

(v) CURRENT APPLICATION DATA:

APPLICATION NUMBER: EP 97203781.6

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: other nucleic acid

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

TACCACTACA ATGGATG
17

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 658 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
 (B) LOCATION: 89..716
 (D) OTHER INFORMATION: /label= N
 /note= "N" stands for unknown."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

CGGAGTTACA AGAGGCTACC AAAGAATTCC AGATTAAAGA GGATATTCTT
 GAAACAAAGA 60

TGAAATTCTT ATCAGTTGAA ACTCCTGANA ATGACAGCCA GTTGTCAAAT
 ATCTCCTGTT 120

CGTTTCAAGG TGAGCTCGAA AGTTCCTTAT GAGATACAAA AAGGACAATG
 CACTTATCAC 180

CTTTGAAAAA GGAAGAAGTT GCTCAAAATG TNGTAANGC ATGAGTAAAC
 ATCATGTACA 240

GATAATAAGA TGTAATCTG GAGGTTACGG CCAAAGCCAA GTTCCATTAA
 TATTCAAGGA 300

GTCANGATTC CAGNGTTTAT GCTAGAANGT TTCTAAAAAT GANAATCAAT
 GGTTACTGGA 360

AATTCCTGGA CACATTGCGN TGAAAGATCA AGATGACGAA GACAAACTAA
 GAAGCTGAGC 420

TTTTCAAAAG TCCCGAAANA TGGAAGAGCG GTAGAGGGTG GNACCGCGTG
 NGANCTATGA 480

CAAGACAAGN CCGGGGAAGN TGCAGTCCAT CACGTTTGTN NGAAGATTGG
 ANGTNGGCTG 540

ACCAANGAAT TTTGAAAAAG GAGANGAATT ACCCCTCTTT ANGAGTAANA
 TCAAAACCCT 600

GCCATAANAA GTTNACTGGT TTCNCCCATT ACACAGNANT TACANNTTGA
 NCAANANTAN 660

NCAGGATAAT TTNCAGGGGA ANAATCTNAA GNATGGCAAG NTGACTTCTG
 GACAANGGT 719

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 220 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: unknown
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(2) INFORMATION FOR SEQ ID NO: 5:

(A) LENGTH: 307 amino acids

(C) STRANDEDNESS: un

(D) TOPOLOGY: unknown

(iii) HYPOTHETICAL: NO

Met Glu Ala Asp Lys Asp Asp Thr Gln Gln Ile Leu Lys
Glu His Ser
1 5 10

Ile Thr Lys Lys Asn Ile Gln Leu Lys Lys Glu Ile Gln
Lys Leu Glu 35 40 45

	Pro	Glu	Thr	Lys	Met	Lys	Phe	Leu	Ser	Val	Glu	Thr	Pro
Glu	Asn	Asp											
	65					70					75		

95

Pro Tyr Glu Ile Gln Lys Gly Gln Ala Leu Ile Thr Phe
Glu Lys Glu

305

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 659 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(ix) FEATURE:

- (A) NAME/KEY: misc feature
- (B) LOCATION: 189..657
- (D) OTHER INFORMATION: /label= N
/note= "N" stands for unknown.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

AGCAGGTGCT GCAACAAAG GAGCACACGA TCAACATGGA GGAGTGCCGG
CTGCGGGTGC 60

AGGTCCAGCC CTTGGAGCTG CCCATGGTCA CCACCATCCA GGTGTCCAGC
CAGTTGAGTG 120

GCCGGAGGGT GTTGGTCACT GGATTTCTTG CCAGCCTCAG GCTGAGTGAG
GAGGAGCTGC 180

TGGACAANCT ANAGATCTTC TTTGGCAAGA CTAGGAACGG AGGTGGCNAT
GTGGACNTTC 240

GGGANCTACT GCCAGGGANT GTCATGCTGG GGTTTGCTAG GGATGGAGTG
GCTCANCCTC 300

TGTGCCAAAT CGGCCATTTT ACAGTGCCAC TGGGTGGGCA GCANGTCCCT
CTGAGAGTCT 360

CTCCGTATGT GAATGGGGAN ATCCAGANGG CTGANATCAG GTCNCAGCCA
NTTCCCCGCT 420

CGGTACTGGT GCTCAACATT CCTGATACTT TGGATGGCCC GGAGCTGCAT
GACGTCCTGG 480

ANATCCACTT CCAGAANCCC ACCCGCGGGG GCGGAGATGT AAGACGCCCT
GACAGTCGTA 540

CCCCAAGGAC AACAGGGCCT AACAGTCTTC ACCTCCTGAA TCAAGGCTAN
GGGCCTCCCC 600

CTTCTCATCC TCCCCACCCC CCCCGCCAAA GGTTCTCAAN ACTGGGCCTG
GGCTTTNTG 659

(2) INFORMATION FOR SEQ ID NO: 7:

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(ix) FEATURE:

- (A) NAME/KEY: misc feature
- (B) LOCATION: 182..626
- (D) OTHER INFORMATION: /label= N
/note= "N" stands for unknown."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

GGATCCACTG CCCTCTGCTT GCGGGCTCTG CTCTGATCAC CTTTGATGAC
CCCAAAGTGG 60

CTGAGCAGGT GCTGCAACAA AAGGAGCACA CGATCAACAT GGAGGAGTGC
CGGCTGCGGG 120

TGCAGGTCCA GCCCTTGGAG CTGCCCATGG TCACCACCAT CCAGGTGATG
GTGTCCAGCC 180

ANTTGAGTGG CCGGAGGGTG TTGGTCACTG GATTTCTGTC CAGCCTCAGG
CTGANTGAGG 240

AGGAGCTGCT GGACAAGCTA TGAGATCTTC TTTGGCAANA CTANGAACGG
ANGTGGCGAT 300

GTGGACGTTT GGGAGCTACT GCCAGGGAGT GTCATGCTGG GGTTCGCTAC
GGATGGAGTG 360

GCTCAGCGTC TGTGCCAAAT CGGCCAGTTC ACAAGTGCCA CTGGGTGGGC
AGCAAGTCCC 420

TCTGAGAGTC TCTCCGTATG TGANTGGNGA GATCAGAATG CTGANATTAA
GTCGCATCCA 480

ATTCCTCGCT CNGGTACTGG TGCTCANNAT CCTGANATCT TGGATTGGCC
CCNGANTNCA 540

TGANATCTGG NAGATTCAAT TNCANAAGTC CANCCNNCNG NGNCGGGAAG
TANANGCCCG 600

ANANTTCNTN NCNTANGGNC AGCANNGCCT G
631

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 138 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

10 Met Ser Ala Pro Leu Asp Ala Ala Leu His Ala Leu Gln
 Glu Glu Gln 1 5 10
 15 Ala Arg Leu Lys Met Arg Leu Trp Asp Leu Gln Gln Leu
 Arg Lys Glu 20 25
 30 Leu Gly Asp Ser Pro Lys Asp Lys Val Pro Phe Ser Val
 20 Pro Lys Ile 35 40 45
 Pro Leu Val Phe Arg Gly His Thr Gln Gln Asp Pro Glu
 Val Pro Lys 50 55 60
 25 Ser Leu Val Ser Asn Leu Arg Ile His Cys Pro Leu Leu
 Ala Gly Ser 65 70 75
 80 Ala Leu Ile Thr Phe Asp Asp Pro Lys Val Ala Glu Gln
 Val Leu Gln 85 90
 95 Gln Lys Glu His Thr Ile Asn Met Glu Glu Cys Arg Leu
 35 Arg Val Gln 100 105
 110 Val Gln Pro Leu Glu Leu Pro Met Val Thr Thr Ile Gln
 40 Val Ser Ser 115 120 125
 Gln Leu Ser Gly Arg Arg Val Leu Val Thr Gly Phe Pro
 Ala Ser Leu 130 135 140
 45 Arg Leu Ser Glu Glu Glu Leu Leu Asp Lys Leu Glu Ile
 Phe Phe Gly 145 150 155
 160 Lys Thr Arg Asn Gly Gly Gly Asp Val Asp Val Arg Glu
 50 Leu Leu Pro
 55

EP 0 921 192 A1

		20		25
30				
5	His Thr Ile Asn Met Glu Glu Cys Arg Leu Arg Val Gln			
	Val Gln Pro	35	40	45
	Leu Glu Leu Pro Met Val Thr Thr Ile Gln Val Met Val			
10	Ser Ser Xaa	50	55	60
	Leu Ser Gly Arg Arg Val Leu Val Thr Gly Phe Pro Ala			
	Ser Leu Arg	65	70	75
15	80			
	Leu Xaa Glu Glu Glu Leu Leu Asp Lys Leu Asp Leu Leu			
	Trp Gln Xaa	85	90	
	95			
20	Xaa Glu Arg Xaa Trp Arg Cys Gly Arg Ser Gly Ala Thr			
	Ala Arg Glu	100	105	
	110			
25	Cys His Ala Gly Val Cys Tyr Gly Trp Ser Gly Ser Ala			
	Ser Val Pro	115	120	125
	Asn Arg Pro Val His Lys Cys His Trp Val Gly Ser Lys			
30	Ser Leu Glu	130	135	140
	Ser Leu Arg Met Xaa Xaa Arg Ser Glu Cys Xaa Val Ala			
	Ser Asn Ser	145	150	155
35	160			
	Ser Leu Xaa Tyr Trp Cys Ser Xaa Ser Xaa Leu Gly Leu			
	Ala Pro Xaa	165	170	
	175			
40	Xaa Met Xaa Ser Gly Arg Phe Asn Xaa Xaa Ser Pro Xaa			
	Xaa Xaa Xaa	180	185	
	190			
45	Gly Lys Xaa Xaa Pro Xaa Xaa Ser Xaa Xaa Xaa Xaa Ser			
	Xaa Ala	195	200	205

(2) INFORMATION FOR SEQ ID NO: 12:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 647 amino acids
 - (B) TYPE: amino acid

Asp Leu Ser Leu Lys Ile Pro Glu Ile Ser Ile Gln Asp
Met Thr Ala

165

170

175

Gln Val Thr Ser Pro Ser Gly Lys Thr His Glu Ala Glu
Ile Val Glu

180

185

190

Gly Glu Asn His Thr Tyr Cys Ile Arg Phe Val Pro Ala
Glu Met Gly

195

200

205

Thr His Thr Val Ser Val Lys Tyr Lys Gly Gln His Val
Pro Gly Ser

210

215

220

Pro Phe Gln Phe Thr Val Gly Pro Leu Gly Glu Gly Gly
Ala His Lys

225

230

235

Val Arg Ala Gly Gly Pro Gly Leu Glu Arg Ala Glu Ala
Gly Val Pro

245

250

255

Ala Glu Phe Ser Ile Trp Thr Arg Glu Ala Gly Ala Gly
Gly Leu Ala

260

265

270

Ile Ala Val Glu Gly Pro Ser Lys Ala Glu Ile Ser Phe
Glu Asp Arg

275

280

285

Lys Asp Gly Ser Cys Gly Val Ala Tyr Val Val Gln Glu
Pro Gly Asp

290

295

300

Tyr Glu Val Ser Val Lys Phe Asn Glu Glu His Ile Pro
Asp Ser Pro

305

310

315

Phe Val Val Pro Val Ala Ser Pro Ser Gly Asp Ala Arg
Arg Leu Thr

325

330

335

Val Ser Ser Leu Gln Glu Ser Gly Leu Lys Val Asn Gln
Pro Ala Ser

340

345

350

Phe Ala Val Ser Leu Asn Gly Ala Lys Gly Ala Ile Asp
Ala Lys Val

Val Ala Lys Gly Leu Gly Leu Ser Lys Ala Tyr Val Gly
Gln Lys Ser

565

570

575

Ser Phe Thr Val Asp Cys Ser Lys Ala Gly Asn Asn Met
Leu Leu Val

580

585

590

Gly Val His Gly Pro Arg Thr Pro Cys Glu Glu Ile Leu
Val Lys His

595

600

605

Val Gly Ser Arg Leu Tyr Ser Val Ser Tyr Leu Leu Lys
Asp Lys Gly

610

615

620

Glu Tyr Thr Leu Val Val Lys Trp Gly His Glu His Ile
Pro Gly Ser

625

630

635

640

Pro Tyr Arg Val Val Val Pro

645

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 213 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: unknown

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

His Glu Gly Arg Gly Val Thr Gly Asn Pro Ala Glu Phe
Val Val Asn

1

5

10

15

Thr Ser Asn Ala Gly Ala Gly Ala Leu Ser Val Thr Ile
Asp Gly Pro

20

25

30

Ser Lys Val Lys Met Asp Cys Gln Glu Cys Pro Glu Gly
Tyr Arg Val

35

40

45

(iii) HYPOTHETICAL: NO

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

10

His Glu Gly Arg Pro Thr Glu Pro Gly Asn Tyr Ile Ile
Asn Ile Lys

1 5 10
15

15

Phe Ala Asp Gln His Val Pro Gly Ser Pro Phe Ser Val
Lys Val Thr

20 25
30

20

Gly Glu Gly Arg Val Lys Glu Ser Ile Thr Arg Arg Arg
Arg Ala Pro

35 40 45

Ser Val Ala Asn Val Gly Ser His Cys Asp Leu Ser Leu
Lys Ile Pro

50 55 60

25

Glu Ile Ser Ile Gln Asp Met Thr Ala Gln Val Thr Ser
Pro Ser Gly

65 70 75
80

30

Lys Thr His Glu Ala Glu Ile Val Glu Gly Glu Asn His
Thr Tyr Cys

85 90
95

35

Ile Arg Phe Val Pro Ala Glu Met Gly Thr His Thr Val
Ser Val Lys

100 105
110

40

Tyr Lys Gly Gln His Val Pro Gly Ser Pro Phe Gln Phe
Thr Val Gly

115 120 125

Pro Leu Gly Glu Gly Gly Ala His Xaa Val Arg Ala Gly
Gly Pro Gly

130 135 140

45

Leu Xaa Lys Ser Ser Trp Ser Ala Ser Arg Ile Gln Tyr
Leu Gly Pro

145 150 155

160

50

Gly Lys Leu Val Leu Glu Ala Trp Pro Leu Leu Ser Xaa
Ala Pro Ala

165 170
175

55

induction of apoptosis in a population of cells related to a pathological condition.

- 5 14. An Nmi/Hou-like proteinaceous substance having at least a functional and/or specific part of the sequence of figure 3 or being encoded by a functional and/or specific part of the sequence of figure 1 or figure 2 or being at least 60, preferably 70, preferably 80% homologous to at least a functional and/or specific part of the sequence of figure 3 or being at least 60, preferably 70, preferably 80% homologous to a protein encoded by at least a functional and/or specific part of the sequence of figure 1 or figure 2.
- 10 15. A recombinant or isolated proteinaceous substance comprising at least a functional part of a member of the family of Nmi/Hou-like proteins or at least a functional part of a member of the family of Hou-like proteins for use in the induction of apoptosis in a population of cells related to a pathological condition.
- 15 16. An IFP35-like proteinaceous substance having at least a functional and/or specific part of the sequence of figure 6 or 7 or being encoded by a functional and/or specific part of the sequence of figure 4 or figure 5 or being at least 60, preferably 70, preferably 80% homologous to at least a functional and/or specific part of the sequence of figure 6 or 7 or being at least 60, preferably 70, preferably 80% homologous to a protein encoded by at least a functional and/or specific part of the sequence of figure 4 or figure 5.
- 20 17. A method for inducing apoptosis in cells comprising providing said cells with Nmi/Hou-like protein activity and/or IFP-35-like activity together with apoptin-like activity.
18. Use of apoptin to find proteinaceous substances associated with apoptosis.

25

30

35

40

45

50

55

CGGAGTTACAAGAGGCTACCAAAGAATTCCAGATTAAAGAGGATATTCCTGAAACAAAGATGAAA
TTCTTATCAGTTGAAACTCCTGANAATGACAGCCAGTTGTCAAATATCTCCTGTTTCGTTTCAAGG
TGAGCTCGAAAGTTCCTTATGAGATACAAAAGGACAATGCACTTATCACCTTTGAAAAGGAAG
AAGTTGCTCAAATGTGNGTAANGCATGAGTAAACATCATGTACAGATAATAAGATGTAAATCTG
GAGGTTACGGCCAAAGCCAAGTTCATTAAATATCAAGGAGTCANGATTCCAGNGTTTATGCTAG
AANGTTTCTAAAAATGANAATCAATGGTTACTGGAAATTCCTGGACACATTGCGNTGAAAGATCA
AGATGACGAAGACAACTAAGAAGCTGAGCTTTTCAAAGTCCCGAAANATGGAAGAGCGGTAGA
GGGTGGNACCGCGTGNGANCTATGACAAGACAAGNCCGGGGAAGNTGCAGTCCATCACGTTTGTN
NGAAGATTGGANGTNGGCTGACCAANGAATTTTGAAAAGGAGANGAATTACCCCTCTTTANGAG
TAANATCAAACCCTGCCATAANAAGTTNACTGGTTTCNCCCATTACACAGNAN
TTACANNTTGANCAANANTANNCAGGATAATTTNCAGGGGAANAATCTNAAGNATGGCAAGNTGA
CTTCTGGACAANGGT

Figure 2

Hou c17/#2

AGCAGGTGCTGCAACAAAAGGAGCACACGATCAACATGGAGGAGTGCCGGCTGCGGGTGCAGGTC
 CAGCCCTTGGAGCTGCCCATGGTCACCACCATCCAGGTGTCCAGCCAGTTGAGTGGCCGGAGGGT
 GTTGGTCACTGGATTTCTGCCAGCCTCAGGCTGAGTGAGGAGGAGCTGCTGGACAANCTANAGA
 TCTTCTTTGGCAAGACTAGGAACGGAGGTGGCNATGTGGACNTTCGGGANCTACTGCCAGGGANT
 GTCATGCTGGGGTTTGCTAGGGATGGAGTGGCTCANCCTCTGTGCCAAATCGGCCATTTACAGT
 GCCACTGGGTGGGCAGCANGTCCCTCTGAGAGTCTCTCCGTATGTGAATGGGGANATCCAGANGG
 CTGANATCAGGTCNCAGCCANTTCCCCGCTCGGTACTGGTGCTCAACATTCCTGATATCTTGGAT
 GGCCCGGAGCTGCATGACGTCTTGGANATCCACTTCCAGAANCCACCCGCGGGGGCGGAGATGT
 AAGACGCCCTGACAGTCGTACCCCAAGGACAACAGGGCCTAACAGTCTTCACCTCCTGAATCAAG
 GCTANGGGCCTCCCCCTTCTCATCCTCCCCACCCCCCGCCAAAGGTTCTCAANACTGGGCCTG
 GGCTTTNTG

Figure 4

IFP35 c14/#1

GGATCCACTGCCCTCTGCTTGCGGGCTCTGCTCTGATCACCTTTGATGACCCCAAAGTGGCTGAG
 CAGGTGCTGCAACAAAAGGAGCACACGATCAACATGGAGGAGTGCCGGCTGCGGGTGCAGGTCCA
 GCCCTTGGAGCTGCCCATGGTCACCACCATCCAGGTGATGGTGTCCAGCCANTTGAGTGGCCGGA
 GGGTGTGGTCACTGGATTTCTGCCAGCCTCAGGCTGANTGAGGAGGAGCTGCTGGACAAGCTA
 TGAGATCTTCTTTGGCAANACTANGAACGGANGTGGCGATGTGGACGTTCTGGGAGCTACTGCCAG
 GGAGTGTCTGCTGGGGTTTGCTACGGATGGAGTGGCTCAGCGTCTGTGCCAAATCGGCCAGTTC
 ACAAGTGCCACTGGGTGGGCAGCAAGTCCCTCTGAGAGTCTCTCCGTATGTGANTGGNGAGATCA
 GAATGCTGANATTAAAGTCGCATCCAATTCCTCGCTCNGGTACTGGTGCTCANNATCCTGANATCT
 TGGATTGGCCCCNGANTNCATGANATCTGGNAGATTCAATTNCANAAGTCCANCCNNCNGNGNCG
 GGAAGTANANGCCCGANANTTCNTNNCNTANGGNCAGCANNGCCTG

Figure 6

IFP35 c51/#3

Filamin 1 RLRNGHVGISFVPKATGEHLVHVKNKGQHVASSFIPVVISQSZIGDASRVVSGQQLHKG
c50/#1 1 -----
c57/#2 1 -----

Filamin 61 HTFEPARFIIDTRDAQYGGLSLSLIEGPSKVDINTEDLEDGTCRVTYCPTPEPGNYIINIKF
c50/#1 1 -----
c57/#2 1 -----HEGRPTPEPGNYIINIKF

Filamin 121 ADQHVPGPSFVSVKVTGEGRVKESITRRRRAPSVANVGSHCDLSLKIPEISIQDHTAQVTS
c50/#1 1 -----
c57/#2 18 ADQHVPGPSFVSVKVTGEGRVKESITRRRRAPSVANVGSHCDLSLKIPEISIQDHTAQVTS

Filamin 181 PSCKTHEAEIVEGENHTYTCIRFVPAEMGHTHTSVVKYKGQHVPGSPFQFTVGPLGEGGAHK
c50/#1 1 -----
c57/#2 78 PSCKTHEAEIVEGENHTYTCIRFVPAEMGHTHTSVVKYKGQHVPGSPFQFTVGPLGEGGAHK

Filamin 241 VRAGGPGLPPEGVPEFS.EWTREAGAGLAAVEPEKAEISFEDRDSCGLAYV
c50/#1 1 -----
c57/#2 138 VRAGGPGLPPEGVPEFS.EWTREAGAGLAAVEPEKAEISFEDRDSCGLAYV

Filamin 300 QPFGDYEVSVKPFNEEHIPDSPFVVPVASPSGDARRLTVSSLQESGLKVNQPASFAVSLNG
c50/#1 1 -----
c57/#2 197 XPESD*NXPIQVSTKEHX-----

Filamin 360 AKGAIDAKVHSPSGALEECYVTEIDQDKYAVRFIPRENGVYLIDVKFNGTHIPGSPFKIR
c50/#1 1 -----
c57/#2 214 -----

Filamin 420 VGEFPGHGGDPGLVSAYCAGLEG.GVTGNPAEFVVNTSNAGAGALSVTIDGPSKVKMDCQE
c50/#1 1 -----
c57/#2 214 -----HEGRGVTGNPAEFVVNTSNAGAGALSVTIDGPSKVKMDCQE

Filamin 479 CPEGYRVITYTPMAPGSYLISIKYGGPYHIGGSPFKAKVTGPRLVSNHSLHETSSVFVDSL
c50/#1 42 CPEGYRVITYTPMAPGSYLISIKYGGPYHIGGSPFKAKVTGPRLVSNHSLHETSSVFVDSL
c57/#2 214 -----

Filamin 539 TKATCAPGHGAPGPGPADASKVVAKGLGLSKAYVQKSSFTVDCSKAONNMLLVGVHGP
c50/#1 102 TKATCAPGHGAPGPGPADASKVVAKGLGLSKAYVQKSSFTVDCSKAONNMLLVGVHGP
c57/#2 214 -----

Filamin 599 TPCDEILVKHVGS.RMYSVSYLLKDKGE.YTLVVKWGHHEHIGSFYRVVP-
c50/#1 162 TPCDEILVKARGQPALQRVLTCPKDKGEVHTGGQNGGQYQIPCKPLP.CGCP
c57/#2 214 -----

Figure 8



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 97 20 3781

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
A	ZHUANG S -M ET AL: "APOPTIN, A PROTEIN ENCODED BY CHICKEN ANEMIA VIRUS, INDUCES CELL DEATH IN VARIOUS HUMAN HEMATOLOGIC MALIGNANT CELLS IN VITRO" LEUKEMIA, vol. 9, no. SUPPL. 01, October 1995, pages S118-S120, XP000602147 * the whole document *	1-7,9-15	TECHNICAL FIELDS SEARCHED (Int.Cl.6)
A	ZHUANG S -M ET AL: "APOPTIN, A PROTEIN DERIVED FROM CHICKEN ANEMIA VIRUS, INDUCES P53- INDEPENDENT APOPTOSIS IN HUMAN OSTEOSARCOMA CELLS" CANCER RESEARCH, vol. 55, no. 3, 1 February 1995, pages 486-489, XP000602162 * the whole document *	1-7,9-15	
T	DE 196 28 894 A (HAGENMAIER HANS PAUL) 22 January 1998 * claims 1-14,16,17 *	1,13	

EPO FORM 1503 03.93 (P04C10)